arXiv:1407.4825v1 [math.RA] 12 Jul 2014

HOCHSCHILD COHOMOLOGICAL DIMENSION IS NOT UPPER-SEMICONTINUOUS

ANASTASIS KRATSIOS

ABSTRACT. In this paper it is demonstrated that the Hochschild Cohomological dimension of an associative algebra is not an upper-semi continuous function, showing the semicontinuity theorem is no longer valid for noncommutative algebras. This is exhibited by a family of $\mathbb{C}$-algebras parametrised by $\mathbb{C}$ all but one of which has Hochschild cohomological dimension 2 and the other having Hochschild cohomological dimension 1.

All algebras are assumed to be unital and associative. Denote by $HH^n(A,M)$ the n^{th} Hochschild cohomology of A with coefficients in M.

Definition 0.1. *The Hochschild cohomological dimension* of a $\mathbb{C}$-algebra A, denoted $HCDim(A)$ is defined as the largest natural number n for which there exists an (A,A)-bimodule M such that $HH^n(A,M) \ncong 0$. If no such number exists then $HCDim(A) := \infty$ [1].

The Hochschild cohomological dimension of the family of $\mathbb{C}$-algebras:

$$A_a := \mathbb{C} < x, y > /(xay - ayx - x) \tag{0.1}$$

parameterized by $a \in \mathbb{C}$ is the focus of this paper.

Lemma 0.2. $HCDim(A_1) = 2$.

Proof.

(1) The Poincaré-Birkhoff-Witt theorem implies A_1 is isomorphic to the universal enveloping algebra of the 2-dimension complex lie algebra $\mathfrak{g}$ with $\mathbb{C}$-basis $\{x, y\}$ who's Lie bracket relations is described on basis elements as $[x,x] = [y,y] = 0$ and $[x,y] = -[y,x] = x$ [2].

(2) Therefore there is an A_1-projective resolution of A_1 of length 2 [4]:

$$0 \leftarrow A_1 \overset{d^0}{\leftarrow} A_1 \otimes_{\mathbb{C}} \mathbb{C}^2 \overset{d^1}{\leftarrow} A_1 \otimes_{\mathbb{C}} \mathbb{C} \leftarrow 0 \tag{0.2}$$

(3) Hence, there are isomorphisms:

$$HH^1(A_1,A_1) \cong Ext^1_{A_1^e}(A_1,A_1) \cong Ext^1_{A_1}(A_1, Hom_{\mathbb{C}}(A_1,A_1)) \tag{0.3}$$

$$\cong Hom_{A_1}(A_1, Hom_{\mathbb{C}}(A_1,A_1)) \cong Hom_{\mathbb{C}}(A_1,A_1). \tag{0.4}$$

(The isomorphisms in (0.3) are described in [4] (for general k-algebras A) and the isomorphisms (0.4) arise from the calculation of the homology of $Hom_{\mathbb{C}}(-, Hom_{\mathbb{C}}(A_1,A_1))$ via to the A_1-projective resolution (0.2).)

Since A_1 is an infinite dimension $\mathbb{C}$-vector space, $Hom_{\mathbb{C}}(A_1,A_1) \ncong 0$, whence $HH^1(A_1,A_1) \ncong 0$. Therefore, $1 \leq HCDim(A)$.

Date: January 1, 2001 and, in revised form, June 22, 2001.

2000 *Mathematics Subject Classification.* Primary 16E40; Secondary 16E40, 13C15.

Key words and phrases. Differential geometry, algebraic geometry.

(4) The resolution (0.2) implies that for $n > 1$ there are isomorphisms:

$$HH^n(A_1, N) \cong Ext^n_{A_1^e}(A_1, N) \cong Ext^n_{A_1}(A_1, Hom_{\mathbb{C}}(A_1, N))$$

$$\cong Hom_{A_1}(0, Hom_{\mathbb{C}}(A_1, A_1)) \cong 0. \tag{0.5}$$

Therefore, $1 \leq HCDim(A) < 2$.

□

Theorem 0.3. *There exists a family $\mathbb{C}$-algebras A_a, parametrised by $\mathbb{C}$ such that if $a \neq 0$ then $HCDim(A_a) = 2$ and $HCDim(A_0) = 1$.*

Proof.

- $A_0 = \mathbb{C} < x, y > /(x) \cong \mathbb{C} < y > \cong \mathbb{C}[x]$. Since $\mathbb{C}[x]$ is a regular $\mathbb{C}$-algebra of finite type it satisfies Van Den Bergh duality in dimension 1 [3]; together with the Hochschild-Kronstadt-Rosenberg theorem [4] this means there are natural isomorphisms for every $(\mathbb{C}[x], \mathbb{C}[x])$-bimodule M:

$$HH^n(\mathbb{C}[X], M) \cong HH_{1-n}(\mathbb{C}[X], Der_{\mathbb{C}}(\mathbb{C}[x], \mathbb{C}[x]) \otimes_{\mathbb{C}[x]} M)$$

$$\cong HH_{1-n}(\mathbb{C}[X], M). \tag{0.6}$$

  In particular, this implies the Hochschild cohomology vanishes for $n > 1$ and

$$HH^1(\mathbb{C}[X], \mathbb{C}[X]) \cong HH_0(\mathbb{C}[X], \mathbb{C}[x]) \cong Z(\mathbb{C}[X]) = \mathbb{C}[X]. \tag{0.7}$$

  Thus, $HCDim(A_0) = 1$.
- On the other hand if $a \neq 0$, A_a is isomorphic to $A_1 := \mathbb{C}\langle x, y\rangle/(xy - yx - x)$ via the $\mathbb{C}$-algebra isomorphisms $\psi_a : A_a \to A_1$ mapping $y \mapsto ay$ (with inverse $y \mapsto \frac{1}{a}y$).

  Hochschild cohomology is functorial [4], therefore for every (A, A)-bimodule M, for every $n \in \mathbb{N}$ and $a \in \mathbb{C} - \{0\}$ ψ_a induces isomorphisms

$$HH^n(\psi_a, M) : HH^n(A_1, M) \to HH^n(A_a, M). \tag{0.8}$$

  Whence $HCDim(A_a) = HCDim(A_1)$ for all $a \in \mathbb{C} - \{0\}$. Therefore it is enough to compute the Hochschild cohomological dimension of A_1.
- Lemma 0.2 entails $HCDim(A_a) = 2$ for all $a \in \mathbb{C} - \{0\}$. Hence $HCDim(A_a) = 2$ if $a \in \mathbb{C} - \{0\}$.

□

References

1. J. Cuntz, D. Quillen, *Algebra extensions and nonsingularity* Journal of the American Mathematical Society **8** (1995), 251–289.
2. T.Y. Lam, *A first course in noncommutative rings* Springer **131** (2001).
3. M. Van den Bergh, *A relation between Hochschild homology and cohomology for Gorenstein rings* Proceedings of the American Mathematical Society **126** (1998), 1345–1348.
4. C. Weibel, *An introduction to homological algebra* Cambridge Univ. Press **38** (2008).

Department of Mathematics, Université de Montréal, Montréal, Québec
E-mail address: kratsiosa@dms.umontreal.ca